

Involute-Evolute Curves in Galilean Space G_3

A. Z. AZAK, M. AKYİĞİT, S. ERSOY

apirdal@sakarya.edu.tr, makyigit@sakarya.edu.tr, sersoy@sakarya.edu.tr

Department of Mathematics, Faculty of Arts and Sciences,
Sakarya University, Sakarya, 54187 TURKEY

March 17, 2010

Abstract

In this paper, definition of involute-evolute curve couple in Galilean space is given and some well-known theorems for the involute-evolute curves are obtained in 3-dimensional Galilean space.

Mathematics Subject Classification (2010). 53A35, 51M30.

Keywords: Galilean space, Involute-evolute curve.

1 Introduction

C. Huggens discovered involutes while trying to build a more accurate clock, [1]. Later, the relations Frenet apparatus of involute-evolute curve couple in the space $\mathbb{E}^3$ were given in [2]. A. Turgut examined involute-evolute curve couple in $\mathbb{R}^n$, [3]. At the beginning of the twentieth century, Cayley-Klein discussed Galilean geometry which is one of the nine geometries of projective space. After that, the studies with related to the curvature theory were maintained [4, 5, 6] and the properties of the curves in the Galilean space were studied in [7]. In this paper, we define involute-evolute curves couple and give some theorems and conclusions, which are known from the classical differential geometry, in the three dimensional Galilean space G_3 . We hope these results will be helpful to mathematicians who are specialized on mathematical modeling.

2 Preliminaries

The Galilean space G_3 is a Cayley-Klein space equipped with the projective metric of signature $(0,0,+,+)$, as in [8]. The absolute figure of the Galilean Geometry consist of an ordered triple $\{w, f, I\}$, where w is the ideal (absolute) plane, f is the line (absolute line) in w and I is the fixed elliptic involution of points of f , [6]. In the non-homogeneous coordinates the similarity group H_8

has the form

$$\begin{aligned}\bar{x} &= a_{11} + a_{12}x \\ \bar{y} &= a_{21} + a_{22}x + a_{23}y \cos \varphi + a_{23}z \sin \varphi \\ \bar{z} &= a_{31} + a_{32}x - a_{23}y \sin \varphi + a_{23}z \cos \varphi\end{aligned}\tag{2.1}$$

where a_{ij} and φ are real numbers, [5].

In what follows the coefficients a_{12} and a_{23} will play the special role. In particular, for $a_{12} = a_{23} = 1$, (2.1) defines the group $B_6 \subset H_8$ of isometries of Galilean space G_3 .

In G_3 there are four classes of lines:

- i) (proper) non-isotropic lines- they don't meet the absolute line f .
- ii) (proper) isotropic lines- lines that don't belong to the plane w but meet the absolute line f .
- iii) unproper non-isotropic lines-all lines of w but f .
- iv) the absolute line f .

Planes $x = \text{constant}$ are Euclidean and so is the plane w . Other planes are isotropic.

If a curve C of the class C^r ($r \geq 3$) is given by the parametrization

$$r = r(x, y(x), z(x))$$

then x is a Galilean invariant the arc length on C .

The curvature is

$$\kappa = \sqrt{y''(x)^2 + z''(x)^2}$$

and torsion

$$\tau = \frac{1}{\kappa^2} \det(r'(x), r''(x), r'''(x)).$$

The orthonormal trihedron is defined

$$\begin{aligned}T(s) &= \alpha'(s) = (1, y'(s), z'(s)) \\ N(s) &= \frac{1}{\kappa(s)}(0, y''(s), z''(s)) \\ B(s) &= \frac{1}{\kappa(s)}(0, -z''(s), y''(s)).\end{aligned}$$

The vectors T, N, B are called the vectors of tangent, principal normal and bi-normal line of α , respectively. For their derivatives the following Frenet formulas hold

$$\begin{aligned}T'(s) &= \kappa(s)N(s) \\ N'(s) &= \tau(s)B(s) \\ B'(s) &= -\tau(s)N(s),\end{aligned}$$

[9].

Galilean scalar product can be written as

$$\langle u_1, u_2 \rangle = \begin{cases} x_1 x_2 & , \text{ if } x_1 \neq 0 \vee x_2 \neq 0 \\ y_1 y_2 + z_1 z_2 & , \text{ if } x_1 = 0 \wedge x_2 = 0 \end{cases}$$

where $u_1 = (x_1, y_1, z_1)$ and $u_2 = (x_2, y_2, z_2)$. It leaves invariant the Galilean norm of the vector $u = (x, y, z)$ defined by

$$\|u\| = \begin{cases} x & , x \neq 0 \\ \sqrt{y^2 + z^2} & , x = 0. \end{cases}$$

Let α be a curve given first by

$$\begin{aligned} \alpha : I &\rightarrow G_3, & I &\subset \mathbb{R} \\ t &\rightarrow \alpha(t) = (x(t), y(t), z(t)) \end{aligned} \quad (2.2)$$

where $x(t), y(t), z(t) \in C^3$ (the set of three-times continuously differentiable functions) and t run through a real interval.

3 Involute- Evolute Curves in Galilean Space

In this section, we give a definition of involute-evolute curve and obtain some theorems about these curves in G_3 .

Definition 3.1 *Let α and α^* be two curves in the Galilean space G_3 . The curve α^* is called involute of the curve α if the tangent vector of the curve α at the point $\alpha(s)$ passes through the tangent vector of the curve α^* at the point $\alpha^*(s)$ and*

$$\langle T, T^* \rangle = 0,$$

where $\{T, N, B\}$ and $\{T^*, N^*, B^*\}$ are Frenet frames of α and α^* , respectively. Also, the curve α is called the evolute of the curve α^* .

This definition suffices to the define this curve mate as

$$\alpha^* = \alpha + \lambda T$$

(see Figure 1).




Figure 1. Involute-evolute curves

Theorem 3.1 *Let α and α^* be two curves in the Galilean space G_3 . If the curve α^* is an involute of the curve α , then the distance between the curves α and α^* is constant.*

Proof. From definition of involute-evolute curve couple, we know

$$\alpha^*(s) = \alpha(s) + \lambda(s)T(s) \quad (3.1)$$

Differentiating both sides of the equation (3.1) with respect to s and use the Frenet formulas, we obtain

$$T^*(s) = T(s) + \frac{d\lambda}{ds}T(s) + \lambda(s)\kappa(s)N(s).$$

Since the curve α^* is involute of α , $\langle T, T^* \rangle = 0$.
Then we have

$$\frac{d\lambda}{ds} + 1 = 0. \quad (3.2)$$

From the last equation, we easily get

$$\lambda(s) = c - s \quad (3.3)$$

where c is constant. Thus, the equation (3.1) can be written as

$$\alpha^*(s) - \alpha(s) = (c - s)T(s). \quad (3.4)$$

Taking the norm of the equation (3.4), we reach

$$\|\alpha^*(s) - \alpha(s)\| = |c - s|. \quad (3.5)$$

This completes the proof. ■

Theorem 3.2 *Let α and α^* be two curves in Galilean space G_3 . κ, τ and κ^*, τ^* be the curvature functions of α and α^* , respectively. If α is evolute of $\alpha^*(s)$ then there is a relationship*

$$\kappa^* = \frac{\tau}{(c - s)\kappa},$$

where c is constant and s is arc length parameter of α .

Proof. Let Frenet frames be $\{T, N, B\}$ and $\{T^*, N^*, B^*\}$ at the points $\alpha(s)$ and $\alpha^*(s)$, respectively. Differentiating both sides of equation (3.4) with respect to s and using Frenet formulas, we have following equation

$$T^*(s) \frac{ds^*}{ds} = (c - s)\kappa(s)N(s) \quad (3.6)$$

where s and s^* are the arc length parameter of the curves α and α^* , respectively. Taking the norm of the equation (3.6), we reach

$$T^* = N \quad (3.7)$$

and

$$\frac{ds^*}{ds} = (c - s)\kappa(s). \quad (3.8)$$

By taking the derivative of equation (3.7) and using the Frenet formulas and equation (3.8), we obtain

$$\kappa^* N^* = \frac{\tau}{(c - s)\kappa} B. \quad (3.9)$$

From the last equation, we get

$$\kappa^* = \frac{\tau}{(c - s)\kappa}.$$

■

Theorem 3.3 *Let α be the non-planar evolute of curve α^* , then α is a helix.*

Proof. Under assumption s and s^* are arc length parameter of the curves α and α^* , respectively. We take the derivative of the following equation with respect to s

$$\alpha^*(s) = \alpha(s) + \lambda(s)T(s)$$

we obtain that

$$T^* \frac{ds^*}{ds} = \lambda \kappa N.$$

$\{T^*, N\}$ are linearly dependent. We may define function as

$$f = \langle T, T^* \wedge N^* \rangle$$

and take the derivative of the function f with respect to s , we obtain

$$f' = -\tau \langle T, N^* \rangle. \quad (3.10)$$

From the equation (3.10) and the scalar product in Galilean space, we have

$$f' = 0.$$

That is,

$$f = \text{const.}$$

The velocity vector of the curve α always composes a constant angle with the normal of the plane which consist of α^* . Then the non-planar evolute of the curve α^* is a helix. ■

Theorem 3.4 *Let the curves β and γ be two evolutes of α in the Galilean space G_3 . If the points P_1 and P_2 correspond to the point of α , then the angle $\widehat{P_1 P P_2}$ is constant.*

Proof. Let's assume that the curves β and γ be two evolutes of α (see Figure 2). And let the Frenet vectors of the curves α , β and γ be $\{T, N, B\}$, $\{T^*, N^*, B^*\}$ and $\{\tilde{T}, \tilde{N}, \tilde{B}\}$, respectively.




Figure 2. Evolutes of α curve

Following the same way in the proof of the Theorem 3.3, it is easily seen that $\{T, N^*\}$ and $\{T, \tilde{N}\}$ are linearly dependent.

Thus,

$$\langle T, T^* \rangle = 0 \quad (3.11)$$

and

$$\langle T, \tilde{T} \rangle = 0. \quad (3.12)$$

We define a function f as an angle between tangent vector T^* and $\tilde{T}$, that is,

$$f(s) = \langle T^*, \tilde{T} \rangle \quad (3.13)$$

Then, differentiating equation (3.13) with respect to s , we have

$$f'(s) = \kappa^* \frac{ds^*}{ds} \langle N^*, \tilde{T} \rangle + \tilde{\kappa} \frac{d\tilde{s}}{ds} \langle T^*, \tilde{N} \rangle \quad (3.14)$$

where s , s^* and $\tilde{s}$ are arc length parameter of the curves α , β and γ , respectively. Also κ , κ^* and $\tilde{\kappa}$ are the curvatures of the curves α , β and γ , respectively. Considering the equations (3.11), (3.12) and (3.14), we get

$$f'(s) = 0.$$

This means that

$$f = \text{const.}$$

So, the proof is completed. ■

References

- [1] Boyer, C., *A history of Mathematics*, New York: Wiley, (1968).
- [2] Hacısalihoğlu, H. H., *Diferensiyel Geometri*, Ankara Üniversitesi Fen Fakültesi, (2000).
- [3] Turgut, A. and Erdoğan, E., *Involute-evolute curve couples of higher order in R^n and their horizontal lifts in R^n* , Common. Fac. Sci. Univ. Ank. Series A, 41 (3) (1992) 125-130.
- [4] Röschel, O., *Die Geometrie des Galileischen Raumes*, Habilitationsschrift, Leoben, (1984).
- [5] Pavkovic, B. J. and Kamenarovic, I., *The equiform differential geometry of curves in the Galilean space G_3* , Glasnik Matematicki, 22 (42) (1987), 449-457.
- [6] Sipus, Z. M., *Ruled Weingarten surfaces in the Galilean space*, Periodica Mathematica Hungarica, 56 (2) (2008), 213-225.

- [7] Öğrenmiş, A. O., Ergüt, M. and Bektaş, M., On the helices in the Galilean space G_3 , Iranian J. of Sci. Tec, Trans. A., 31 (2) (2007), 177-181.
- [8] Molnar, E., *The projective interpretation of the eight 3-dimensional homogeneous geometries*, Beitrage zur Algebra und Geometrie Contributions to Algebra and Geometry, 38 (2) (1997), 261-288.
- [9] Kamenarovic, I., *Existence theorems for ruled surfaces in the Galilean space G_3* , Rad Hrvatskeakad. Znan. Umj. Mat. 10 (1991), 183-196.